

Exercise During Pregnancy

There are many benefits to be gained from regular low impact exercise during pregnancy. These include physical benefits (lower blood pressure, reduced risk of gestational diabetes) and the prevention of excessive weight gain, blood pressure complications and improved psychological wellbeing as a result of endorphin (pleasure-promoting hormone) release that can improve overall mood and reduce stress.

In addition to pregnancy-specific benefits, there are significant lifelong benefits of regular exercise for all adults including reduced risk of cardiovascular disease, type 2 diabetes and some cancers.


Before you start an exercise program in pregnancy, speak with your doctor or midwife to make sure that you do not have any health issues that may prevent you from participating in regular exercise during your pregnancy.

If there are no health or pregnancy reasons why you should not exercise, you should be encouraged during your pregnancy to participate in regular aerobic and strengthening exercises.

General considerations for exercise during pregnancy:

- include a gradual warm-up and slow and sustained cool-down with each session
- avoid exercising in high temperatures and humidity, ensure adequate hydration and wear loose-fitting clothing
- avoid activities with the possibility of falling (i.e. horse- riding, skiing) or impact trauma to the abdomen (i.e. certain team sport games)
- perform regular exercises to strengthen the pelvic floor muscles (i.e. Pilates and yoga). Avoid activities that add extra load to the pelvic floor (i.e. jumping, bouncing, weightlifting, kickboxing)
- take care with weight-bearing exercise and activities involving frequent changes in direction (i.e. court sports) due to increased risk of injury and changes in balance
- reduce inactive behaviour: minimise the amount of time spent in prolonged sitting and breaking up long periods of sitting as often as possible
- avoid exercising to the point of exhaustion. It is important to listen to your body to avoid overexerting yourself
- be mindful that some exercises involve positions and movements that may be uncomfortable or harmful

Before exercising when pregnant, consult your doctor, midwife or relevant healthcare professional to ensure you are exercising safely.



How much exercise is recommended?

How often should I exercise?

Aim to be physically active on most, preferably all days of the week. If you are currently inactive or overweight, start with 3 to 4 days per week on non-consecutive days.

How hard should I exercise?

Most women should aim for a 'moderate' intensity. This means you should feel like you are working 'somewhat hard'.

For women with a high level of fitness who are accustomed to regular vigorous exercise, there is no evidence to suggest that vigorous exercise during pregnancy is harmful, provided that you listen to your body and adjust your routine over time. A rating of 15 to 16 (equating to 'hard') may be appropriate. However, athletes should be wary of pushing too hard. Pregnancy is not a time for serious competition or aiming to reach peak lifetime fitness. If you were inactive before pregnancy, start slowly and build up your activity to a moderate level of exercise.

How long should I exercise for?

Aim to accumulate 150 to 300 minutes of moderate intensity physical activity each week. Ideally, this should be achieved by being active on most days of the week for at least 30 minutes at a time.

If you are currently inactive or at an unhealthy weight, start with 15 to 20 minutes and slowly build up to 30 minutes per session. While no evidence exists for an upper limit to exercise duration during pregnancy, it is not advisable to extend exercise duration beyond 60 minutes per session, unless the intensity is relatively light.

What type of exercise should I do?

You should be encouraged to participate in both aerobic and strengthening exercises.

Aerobic exercises

Aerobic exercises involve continuous activities that use large muscle groups and elevate the heart and breathing rates to cause some 'huff and puff'. Common examples include:

- walking (aim for a brisk pace)
- stationary cycling
- swimming and other water-based activities (avoid heated spas and hydrotherapy pools)
- if you are already running regularly prior to your pregnancy, there is
 no scientific evidence to say whether you should continue or not. This
 should be decided on an individual basis and in consultation with your
 doctor or midwife. Listen closely to your body and monitor the intensity
 appropriately



Exercise During Pregnancy



Strengthening exercises

Strengthening exercises should be performed twice per week, on non-consecutive days, covering the main muscle groups of the body.


Resistance can be provided by light weights, body weight or elasticised resistance-bands.

Aim to perform 1 to 2 sets of 12 to 15 repetitions for each exercise. These strengthening exercises should be performed at a 'moderate' intensity (rating of perceived exertion 12 to 14), with slow and steady movements and proper breathing technique. (i.e. exhale on exertion).

Avoid heavy weight-lifting and activities that involve straining or holding the breath. Exercises should not be performed lying flat on the back after the first trimester and walking lunges are best avoided to prevent injury to the pelvic connective tissue.

Warning signs to stop exercise and seek medical attention:

- chest pain
- dizziness, feeling faint or headache
- muscle weakness
- calf pain, swelling or redness
- sudden swelling of the ankles, hands or face
- vaginal bleeding or amniotic fluid loss
- decreased fetal movement
- increased incontinence
- uterine contractions or pain in the lower back, pelvic area or abdomen (potentially indicating preterm labour)







How you might describe your exertion:

None Reading a book, watching television

Very, very light Tying shoes

Very Light Chores like folding clothes that seem to take little effort

Fairly light Walking through the grocery store or other activities

that require some effort but not enough to speed up

your breathing

Somewhat hard Brisk walking or other activities that require moderate

effort and speed your heart rate and breathing but

don't make you out of breath

Hard Bicycling, swimming, or other activities that take

vigorous effort and get the heart pounding and make

breathing very fast

Very hard The highest level of activity you can sustain

Very, very hard A finishing sprint in a race or other burst of activity that

you can't maintain for long



Useful resources

- https://www.health.gov.au/resources/publications/evidence-basedphysical-activity-guidelines-for-pregnant-women?language=en
- https://www.health.gov.au/sites/default/files/documents/2021/05/ physical-activity-and-exercise-during-pregnancy-guidelines-brochureguidelines-for-physical-activity-during-pregnancy.pdf
- https://www.health.gov.au/resources/publications/physical-activityadvice-during-pregnancy?language=en
- https://www.health.govt.nz/your-health/pregnancy-and-kids/pregnancy/helpful-advice-during-pregnancy/being-active-during-pregnancy
- RANZCOG statement on exercise duing pregnancy (C-Obs 62) available at: www.ranzcog.edu.au

DISCLAIMER: This document is intended to be used as a guide of general nature, having regard to general circumstances. The information presented should not be relied on as a substitute for medical advice, independent judgement or proper assessment by a doctor, with consideration of the particular circumstances of each case and individual needs. This document reflects information available at the time of its preparation, but its currency should be determined having regard to other available information. RANZCOG disclaims all liability to users of the information provided.

The Royal Australian and New Zealand College of Obstetricians and Gynaecologists

1 Bowen Crescent,

Melbourne, VIC 3004, Australia

Phone: +61 3 9417 1699 Email: ranzcog@ranzcog.edu.au Web: ranzcog.edu.au

